



REC'D 12 SEP 2003	
WIPO	PCT

Kongeriget Danmark

Best Available Copy

Patent application No.: PA 2002 01189
 Date of filing: 7 August 2002
 Applicant: Bavarian Nordic A/S
 (Name and address) Vesterbrogade 149
 1620 Copenhagen V
 Denmark

Title: Vaccinia virus host range genes to increase the titer of avipoxviruses

IPC: C 12 N 15/863

This is to certify that the attached documents are exact copies of the above mentioned patent application as originally filed.



Patent- og Varemærkestyrelsen
 Økonomi- og Erhvervsministeriet

07 July 2003

Ase Damm
 Ase Damm

PRIORITY DOCUMENT
 SUBMITTED OR TRANSMITTED IN
 COMPLIANCE WITH
 RULE 17.1(a) OR (b)

- 7 AUG. 2002

Vaccinia virus host range genes to increase the titer of Avipoxviruses

5 The invention concerns an Avipoxvirus comprising in the viral genome a
Vaccinia virus host range gene or a homologue of said host range gene.
The invention further relates to cells, preferably avian cells, comprising a
Vaccinia virus host range gene or a homologue of said host range gene.
Moreover the invention concerns the use of a Vaccinia virus host range
10 gene or an homologue thereof to increase the titer of avipoxviruses
produced from cells after infection of said cells with the avipoxvirus,
wherein the host range gene is expressed in said cells.

15

Background of the invention

The poxviridae comprise a large family of complex DNA viruses that
replicate in the cytoplasm of vertebrate and invertebrate cells. The
family of poxviridae can be divided into the subfamily chordopoxvirinae
20 (vertebrate poxviruses) and entomopoxvirinae (insect poxviruses).

The chordopoxvirinae comprise several poxvirus species that can be
used as vectors to express exogenous DNA segments encoding antigens
against which an immune response is to be induced. Examples for
25 poxviruses that can be used as live vaccines are Vaccinia virus and
avipoxviruses, such as the canarypoxvirus and the fowlpoxvirus.

The use of Vaccinia viruses to engineer viral vectors for recombinant gene expression and for the potential use as recombinant live vaccines has been disclosed in numerous publications (see e.g. Mackett, M., Smith, G.L. and Moss, B. [1982] P.N.A.S. USA 79, 7415-7419; Smith, G.L.,
5 Mackett, M. and Moss, B. [1984] Biotechnology and Genetic Engineering Reviews 2, 383-407). To construct recombinant Vaccinia viruses, DNA sequences (genes), which code for foreign antigens are introduced into the genome of the Vaccinia virus under the regulation of suitable poxvirus promoters. If the gene is integrated at a site in the viral DNA which is
10 non-essential for the life cycle of the virus, the recombinant Vaccinia virus remains infectious. After infection the recombinant virus expresses the integrated DNA sequence (EP 83286 and EP 110385). The recombinant Vaccinia viruses prepared in this way can be used, on the one hand, as live vaccines for the prophylaxis of infectious diseases, and on the other
15 hand, for the preparation of heterologous proteins in eukaryotic cells.

The use of Vaccinia virus as vector for the development of recombinant live vaccines has been affected by safety concerns and regulations. Most of the recombinant Vaccinia viruses described in the literature are
20 based on the Western Reserve strain of Vaccinia virus. It is known that this strain has a high neurovirulence and is thus poorly suited for use in humans and animals (Morita et al., Vaccine 5, 65-70 [1987]). On the other hand the Modified Vaccinia virus Ankara (MVA) is known to be exceptionally safe. MVA has been generated by long-term serial
25 passages of the Ankara strain of Vaccinia virus (CVA) on chicken embryo fibroblasts (for review see Mayr, A., Hochstein-Mintzel, V. and Stickl, H. [1975] Infection 3, 6-14; Swiss Patent No. 568392). MVA is distinguished by its great attenuation that is to say by diminished virulence or infectiosity while maintaining good immunogenicity.

Recombinant MVA useful as vaccines have already been constructed and used in clinical trials. WO 98/13500 discloses a recombinant MVA containing and capable of expressing one or more DNA sequences encoding dengue virus antigens. The foreign DNA sequences were recombined into the viral DNA at the site of a naturally occurring deletion in the MVA genome.

Another approach towards the generation of safe and effective poxvirus vaccines utilizes avipoxviruses, e.g. canarypoxvirus and fowlpoxvirus, to express antigens to induce an immune response (US 6,340,462). Avipoxviruses are naturally host-restricted and productively replicate only in avian species and cells (Taylor et al., Vaccine 1995, 13: 539-549). If human cells are infected with an avipoxvirus, heterologous genes are expressed from the viral genome. However, the avipoxvirus does not replicate in the human cells and there is, thus, no risk that the human being is harmed by productive virus replication. Various recombinant avipoxviruses have been constructed that express e.g. lentiviral gene products (US 5,766,598), cytokines and/or tumor-associated antigens (US 5,833,975) or rabies G glycoprotein (Taylor et al., Vaccine 1995, 13: 539-549). A recombinant canarypox virus expressing the four HIV genes gag, pol, env and nef has already been used in clinical trials (Peters, B.S., Vaccine 2002, 20: 688-705).

Since avipoxviruses productively replicate only in avian cells, these cells have to be used for the amplification of the virus and for the generation of recombinant viruses. Unfortunately, the titers of avipoxviruses obtained with avian cells are relatively low when compared to other poxviruses and it is, thus, more difficult to produce larger amounts of (recombinant) avipoxviruses in an industrial scale.

Object of the invention

5 It is the object of the present invention to provide means allowing the production of avipoxviruses, in particular recombinant avipoxviruses, at higher titers allowing the production of larger amounts of virus, in particular in an industrial scale.

10

Detailed description of the invention

15 This object has been achieved by expressing vaccinia virus host range genes in cells productively infected with an avipoxvirus. The expression of these vaccinia virus genes leads to an increase of the avipoxvirus titer produced from the infected cells. As it will be shown in more detail in the example section for a specific embodiment of the invention recombinant avipoxviruses, in particular a canarypoxvirus, expressing a
20 Vaccinia virus host range gene, in particular the Vaccinia virus gene C7L, show a 10 fold increase of the viral titer on avian cells, in particular on Chicken Embryo Fibroblasts (CEF-cells) compared to the Avipoxvirus lacking the Vaccinia virus host range gene. Although the host range gene is expressed from the recombinant avipoxvirus, the growth on
25 human cell lines is not affected, i.e. the Avipoxvirus expressing the Vaccinia virus host range gene is as attenuated as the Avipoxvirus not expressing the host range gene.

According to a preferred embodiment the invention concerns avipoxviruses comprising in the viral genome a Vaccinia virus host range gene or a homologue of said host range gene.

5 The term "avipoxvirus" refers to any avipoxvirus, such as Fowlpoxvirus, Canarypoxvirus, Uncopoxvirus, Mynahpoxvirus, Pigeonpoxvirus, Psittacinepoxvirus, Quailpoxvirus, Peacockpoxvirus, Penguinpoxvirus, Sparrowpoxvirus, Starlingpoxvirus and Turkeypoxvirus. Preferred avipoxviruses are Canarypoxvirus and Fowlpoxvirus.

10

An example for a canarypox virus is strain Rentschler. A plaque purified Canarypox strain termed ALVAC (US 5,766,598) was deposited under the terms of the Budapest treaty with the American Type Culture Collection (ATCC), accession number VR-2547. Another Canarypox
15 strain is the commercial canarypox vaccine strain designated LF2 CEP 524 24 10 75, available from Institute Merieux, Inc.

20

Examples of a Fowlpox virus are strains FP-1, FP-5 and TROVAC (US 5,766,598). FP-1 is a Duvette strain modified to be used as a vaccine in one-day old chickens. The strain is a commercial fowlpox virus vaccine strain designated O DCEP 25/CEP67/2309 October 1980 and is available from Institute Merieux, Inc. FP-5 is a commercial fowlpox virus vaccine strain of chicken embryo origin available from American Scientific Laboratories (Division of Schering Corp.) Madison, Wisconsin,
25 United States Veterinary License No. 165, serial No. 30321.

The Vaccinia virus host range gene comprised in the viral genome of the avipoxvirus can be any host range gene. The term "Vaccinia virus host range gene" refers to a gene encoding a gene product, which is

necessary that a Vaccinia virus is able to replicate on cells of species on which the virus does not replicate in the absence of the functional host range gene. If the respective host range gene is deleted the viral replication may be restricted to cells from only one animal species. By way of example reference is made to the Vaccinia virus genes K1L, C7L and E3L. It has been shown that the expression of either K1L or C7L allows vaccinia virus replication in human MRC-5 cells; the E3L gene was shown to be required for Vaccinia virus replication in monkey Vero and human HeLa cells (Wyatt et al., Virology 1998, 251: 334-342).

The term "Vaccinia virus host range gene for human cells" refers to genes that are required for the replication of the Vaccinia virus in human cells.

Examples for Vaccinia virus host range genes are the genes C18L, C17L, C7L, K1L, E3L, B4R, B23R and B24R according to the nomenclature as used in Johnson et al., Virology 1993, 196: 381-401 and genes (CHO)hr and SPI-1 as specified in Wyatt et al., Virology 1998, 251: 334-342.

Preferred host range genes are the host range genes for human cells, e.g. E3L, K1L and C7L. Most preferred is C7L. The sequence of the C7L gene of MVA with regulatory sequences is shown in Figure 4 and as SEQ ID: No 1

The term "homologue of a host range gene" refers to a gene having a homology of at least 50%, preferably at least 70%, more preferably of at least 80%, most preferably of at least 90% in the coding part of the nucleotide sequence, wherein the "homologue of the host range gene" has the biological function of a host range gene. The biological function and definition of a host range gene is given above. Specific tests how to

determine whether a gene has the biological function of a host range gene are known to the person skilled in the art. In particular reference is made to Wyatt et al., Virology 1998, 251: 334-342, Perkus et al., Virology 1990, 179: 276-286 and Gillard et al., J. Virol. 1985, 53: 316-318.

According to the present invention the Vaccinia virus host range gene comprised in the viral genome of the Avipoxvirus is a functional gene. The term "functional gene" as used in the present application is to be interpreted in that the host range gene comprises regulatory elements that are functional in cells productively infected with an Avipoxvirus and that allow the generation of the functional gene product of the host range gene in said Avipoxvirus infected cells. Thus, the Vaccinia virus host range gene is expressed in the cells.

The term "cells, productively infected with an Avipoxvirus" refers to cells that allow the propagation of the Avipoxvirus and/or the generation of recombinant Avipoxviruses. The cells are preferably avian cells, most preferably CEF cells. Other preferred cells are the quail fibroblast cell line QT-35 (Cowen, B.S. and Braune, M.O., Avian Dis. 1988; 32: 282-297; Schnitzlein, W.M. et al., Virus Res. 1988; 10: 65-76) or canary embryo cells (Wurtz, S., Bonnet-Piro, E. and Barban, V., Poster P45 and Wurtz, S. and Barban, V., Poster P93, XIIIth International Poxvirus and Iridovirus Symposium, Montpellier, France, September 2-6, 2000). Further, it is obvious for a person skilled in the art to try whether other, commercially available Avian cell lines are suitable for the propagation of Avipoxviruses. Examples for such cell lines obtainable at the American Type Culture Collection (ATCC) are the quail fibroblast cell lines QT6 (ATCC CRL-1708), QM7 (ATCC CRL-1962), QNR/D (ATCC CRL-

2532), the quail cell line QNR/K2 (ATCC CRL-2533), the duck fibroblast cell line Duck embryo (ATCC CCL-141), the turkey lymphoblast cell line MDTC-PR19 (ATCC CRL-8135), the chicken fibroblast cell lines SL-29 (ATCC CRL-1590) and UMNSAH/DF-1 (ATCC CRL-12203) and the
5 chicken lymphoblast cell lines DT40 (ATCC CRL-2111) and DT95 (ATCC CRL-2112).

The regulatory elements comprise inter alia suitable promoter/enhancer and termination signals that are known to the person skilled in the art
10 as being active in avipoxvirus infected avian cells. Examples for such promoter/enhancer elements are the Vaccinia virus promoters P_{7.5}, P_{H5}, P₁₁, a synthetic strong promoter P_{syn} (see Genetically engineered viruses, edited by Ring, C.J.A. and Blair, E.D., Bios Scientific Publishers Ltd., 2001, Oxford, UK, ISBN 1 85996 103 7, chapter Vaccinia virus
15 promoters, starting on page 110; Amano, H. et al., Virology 1999, 256: 280-290) as well as the autologous promoters of the Vaccinia virus host range genes.

Fang, Z.-Y. et al., (Virology 2001, 291: 272-284) discloses a
20 recombinant canarypoxvirus that comprises in the viral genome three expression cassettes for HIV gag-pro, gp120/TM and a Nef/Pol poly-epitope string and an expression cassette of the E3L gene. The viral genome of said recombinant further comprises the Vaccinia Virus K3L gene. Due to the presence of the vaccinia virus E3L and K3L genes the
25 apoptosis in infected HeLa cells was significantly reduced and the antigen production in the infected cells was enhanced. Fang et al., does not disclose that the Vaccinia virus host range gene is implicated in any increase of the titer of the recombinant avipoxvirus. Thus, the avipoxvirus according to the present invention is an avipoxvirus

comprising in the viral genome a Vaccinia virus host range gene or a
homologue of said host range gene, with the proviso that the host range
gene is not the E3L gene if the avipoxvirus is a recombinant
canarypoxvirus comprising in the viral genome expression cassettes for
5 (I) HIV gag-pro, (II) gp120/TM and a (III) Nef/Pol poly-epitope string as
well as for (IV) the Vaccinia virus K3L gene. According to a further
alternative embodiment the invention concerns an avipoxvirus
comprising in the viral genome a Vaccinia virus host range gene or a
homologue of said host range gene, with the proviso that the host range
10 gene is not the E3L gene if the avipoxvirus is a recombinant
canarypoxvirus comprising in the viral genome the Vaccinia virus K3L
gene.

The Vaccinia virus host range gene is preferably inserted into a non-
15 essential region of the viral genome, into an intergenic region of the viral
genome or into a deletion site of the viral genome. "Non essential
regions" are regions that are not required for the replication of the viral
genome in avian cells and not needed for the production of infectious
viruses. Non-essential regions are known to the person skilled in the art
20 and are disclosed i.a. in US 5,766,598. The insertion of heterologous
genes into the Canarypox virus thymidine kinase gene has been
disclosed by Amano, H. et al. (Virology 1999, 256: 280-290).

"Intergenic regions" in the viral regions are regions that do not contain
25 coding sequences and preferably no regulatory elements. The location of
intergenic regions is known to the person skilled in the art (see e.g.
Alfonso C.L. et al., J. Virol 2000, 74: 3815-3831). An example for an
insertion into an intergenic region is shown in figure 3 and in the
examples section. Thus, it is a preferred embodiment, in particular for a

Canarypoxvirus, to insert the host range gene into the intergenic region between the Tk gene and the adjacent X gene.

5 A deletion site is the part of the genome of a modified avipoxvirus that is deleted with respect to the parent avipoxvirus. Deletion sites may be generated by using methods known by a person skilled in the art, starting from a wild type avipoxvirus genome.

10 The Avipoxvirus comprising a Vaccinia virus host range gene may be a wild-type virus comprising as only heterologous gene the Vaccinia virus host range gene, an attenuated virus comprising as only heterologous gene the Vaccinia virus host range gene or a recombinant Avipoxvirus, i.e. a wild-type or attenuated virus comprising further heterologous genes in addition to the Vaccinia virus host range gene.

15 An "attenuated virus" is a virus originating from a pathogenic virus but that upon infection of the host organism leads to a lower mortality and/or morbidity compared to the non-attenuated parent virus. Examples of attenuated poxviruses are known to the person skilled in the art. Examples for attenuated Avipoxvirus strains are i.a. FP-1, 20 ALVAC or TROVAC.

The term "recombinant virus" refers to any virus that comprises in addition to the vaccinia virus host range gene an additional 25 heterologous nucleic acid that is not naturally part of the viral genome. A heterologous gene can be, e.g., a therapeutic gene, a gene coding for a peptide comprising at least one epitope to induce an immune response, an antisense expression cassette or a ribozyme gene.

Thus, according to a preferred embodiment the invention concerns Avipoxviruses comprising in the viral genome at least one heterologous nucleic acid sequence in addition to the sequence encoding the Vaccinia virus host range gene, wherein the additional heterologous nucleic acid sequence is preferably selected from a sequence coding for at least one antigen, antigenic epitope, and/or a therapeutic compound.

In a preferred embodiment the present invention concerns the avipoxviruses according to the present invention as a vaccine. A „vaccine“ is a compound, i.e. a vector or a virus that induces a specific immune response.

The heterologous nucleic acids are preferably inserted into the preferred insertion sites of the viral genome as explained above for the Vaccinia virus host range genes. Thus, preferred insertion sites for heterologous nucleic acids are i.a. intergenic regions of the viral genome, deletion sites and non-essential regions.

If the Avipoxvirus is a non-recombinant virus, i.e. an avipoxvirus that does not contain in the viral genome heterologous genes other than the Vaccinia virus host range gene, the Avipoxvirus can be used to vaccinate against avian poxvirus infections. This is of significant importance in the veterinary field, e.g. for the vaccination of poultry. In this case it is preferred to use an attenuated Avipoxvirus. If the Avipoxvirus is a recombinant virus, i.e. an avipoxvirus that contains in the viral genome heterologous genes other than the Vaccinia virus host range gene, the Avipoxvirus can be used to vaccinate against avian poxvirus infections and/or to induce an immune response against the peptide/protein that is encoded by the additional heterologous nucleic

acid. This embodiment is of particular importance if a recombinant Avipoxvirus is used for the vaccination of mammals, in particular humans. In this case the additional heterologous sequence may express antigens against which it is intended to induce an immune response.

5 Examples for such antigens are i.a. tumour antigens, antigens derived from infectious agents such as viruses, bacteria, fungi, synthetic polypeptide strings and so on.

10 The vaccination is made by administering an Avipoxvirus according to the present invention to an animal, including an human. The mode of administration, the dose and the number of administrations can be optimized by those skilled in the art in a known manner. Most preferred for poxvirus vectors is subcutaneous or intramuscular administration.

15 For the preparation of a vaccine, the virus according to the invention is converted into a physiologically acceptable form. This can be done based on the experience in the preparation of poxvirus vaccines used for vaccination against smallpox (as described by Stickl, H. *et al.* [1974] Dtsch. med. Wschr. 99, 2386-2392). For example, the purified virus is
20 stored at -80°C with a titer of 5×10^8 TCID₅₀/ml formulated in about 10mM Tris, 140 mM NaCl pH 7.4. For the preparation of vaccine shots, e.g., 10^2 - 10^8 particles of the virus are lyophilized in phosphate-buffered saline (PBS) in the presence of 2% peptone and 1% human albumin in an ampoule, preferably a glass ampoule. Alternatively, the vaccine shots
25 can be produced by stepwise freeze-drying of the virus in a formulation. This formulation can contain additional additives such as mannitol, dextran, sugar, glycine, lactose or polyvinylpyrrolidone or other additives such as antioxidants or inert gas, stabilizers or recombinant proteins (e.g. human serum albumin) suitable for *in vivo* administration.

The glass ampoule is then sealed and can be stored between 4°C and room temperature for several months. However, as long as no need exists the ampoule is stored preferably at temperatures below -20°C. For vaccination the lyophilisate can be dissolved in 0.1 to 0.5 ml of an aqueous solution, preferably physiological saline or Tris buffer, and is administered either systemically or locally, i.e. by parenterally, intramuscularly or any other path of administration known to the skilled practitioner.

According to a related embodiment the invention concerns a method for affecting, preferably inducing an immunological response in a living animal body, including a human, comprising administering the avipoxvirus according to the present invention, the pharmaceutical composition and/or the vaccine according to the present invention to the animal or human to be treated. According to a preferred embodiment the animal may be immuno-compromised. In immuno-compromised animals it is preferred to use severely attenuated virus strains in order to assure that the animal is not overwhelmed by productive virus replication. This may be of particular relevance if the animal is a natural host for the virus, which is the case in poultry. Since Avipoxviruses do not replicate in humans the Avipoxviruses according to the present invention are particularly safe in human beings even if the used virus strain is not an attenuated strain with respect to the natural host.

25

According to a further embodiment the invention relates to a pharmaceutical composition comprising the avipox virus according to the present invention and a pharmaceutically acceptable carrier, diluent and/or additive. The pharmaceutical composition is in fact a vaccine if

the composition comprises an Avipoxvirus containing in the viral genome a heterologous nucleic acid encoding an antigen against which an immune response is to be induced. However, the heterologous nucleic acid is not restricted to this type of sequences. Instead, the heterologous sequence may also be a suicide gene, such as the herpes simplex virus thymidine kinase gene, a therapeutic gene, such as an antisense RNA gene or ribozyme gene or any other gene having an therapeutic benefit. According to the latter alternatives the avipovirus according to the present invention may be part of a pharmaceutical composition aiming at treating disease and not primarily intending to vaccinate against a disease. If the heterologous gene is a suicide gene the pharmaceutical composition may be administered locally to a tumour, leading to the infection of the tumour cells with the recombinant avipoxvirus. The suicide gene is then expressed in the tumour cells and by administration of the prodrug that corresponds to the respective gene product of the suicide gene (e.g. gancyclovir in the case of the Herpes simplex virus thymidine kinase gene) a selective killing of tumour cells becomes possible.

The pharmaceutical composition and/or the vaccine may generally include one or more pharmaceutical acceptable and/or approved carriers, additives, antibiotics, preservatives, adjuvants, diluents and/or stabilizers. Such auxiliary substances can be water, saline, glycerol, ethanol, wetting or emulsifying agents, pH buffering substances, or the like. Suitable carriers are typically large, slowly metabolized molecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, lipid aggregates, or the like.

According to a preferred embodiment the invention concerns a method for introducing a homologous and/or a heterologous nucleic acid sequence into target cells comprising the infection of the target cells with the avipoxvirus according to the present invention. In the context of this embodiment the terms "heterologous" and "homologous" nucleic acid refer to nucleic acids which are heterologous and homologous, respectively, with respect to the cellular genome. Thus, according to this embodiment a "homologous nucleic acid" is a sequence which is homolog to the cellular genome, such as a cellular gene or a derivative thereof, having a nucleotide sequence homology in the coding region of at least 50%, preferably of at least 70%, more preferably of at least 80%, most preferably of at least 90%. According to this embodiment the term "heterologous nucleic acid" refers to nucleic acids having no homologue in the cellular genome. Examples for such heterologous nucleic acids are viral, bacterial and fungal genes. The target cell may be any cell that can be infected with the virus according to the present invention. Thus, the target cell may be an avian cell, such as CEF cells, or mammalian cells, including human cells. The cell may be a primary cell or a cell line. The target cell can be a cell that is cultivated in vitro (i.e. a cell that is cultivated in culture flasks) or a cell that is part of a living organism. Methods how to infect cells are known to the person skilled in the art.

The invention further concerns a method for producing a peptide and/or protein comprising the infection of a host cell with the avipoxvirus according to the present invention, cultivation of the infected host cell under suitable conditions, and isolation and/or enrichment of the peptide and/or protein expressed from the viral genome. The peptide/protein may be a Avipoxvirus protein/peptide. If the

Avipoxvirus expresses a nucleic acid which is heterologous to the viral genome, the peptide/protein may also be the peptide/protein that is expressed from the heterologous nucleic acid. The host cell type is not critical as long as the cell can be infected with the virus and as long as
5 the protein/peptide to be isolated is produced in said cell from the viral vector. The cell may be a cell in which the virus replicates productively or a cell that does not promote productive replication such as human cells.

10 The invention further concerns a method for producing, in particular amplification of the Avipoxvirus according to the present invention comprising the infection of a host cell with the Avipoxvirus according to the present invention, cultivation of the infected host cell under suitable conditions, and isolation and/or enrichment of the virus produced by
15 said host cell. For amplification of the Avipoxvirus it is necessary to infect cells that allow a reproductive replication of the virus. Such cells are known to the person skilled in the art and include avian cells, i.a. CEF cells. Other suitable cells and cell lines have been disclosed above.

20 The invention further concerns cells infected with the Avipoxvirus according to the present invention. The cells may be cells allowing a productive replication of the Avipoxvirus, such as avian cells, in particular CEF cells or cells that can be infected by the Avipoxvirus but do not promote viral replication, such as primary human cells or human
25 cell lines.

Methods for obtaining the Avipoxvirus according to the present invention are known to the person skilled in the art (see e.g. US 5,766,598; US 5,833,975; US 6,340,462). According to a preferred

embodiment such a method may comprise the following steps: In a first step an avipox virus genome and a DNA comprising a host range gene as defined above are introduced into cells in which the virus is able to reproductively replicate. The avipoxvirus genome may already contain heterologous nucleic acids as defined above. The avipoxvirus genome is conveniently introduced into the cell by infection of the cell with the corresponding avipoxvirus. The DNA is preferably introduced in the cell by transfection techniques known to the person skilled in the art. Such techniques include lipofection or Calcium phosphate precipitation. The DNA that is introduced into the cells is preferably capable to specifically recombine with the genomic DNA of the avipoxvirus. To this end the nucleic acid to be inserted into the viral genome is flanked by viral sequences which direct a specific recombination of the nucleic acid into the viral genome. Depending on the type of the flanking viral sequences it is possible to insert the nucleic acid into any part of the viral genome. Preferably the insertion is done into non-essential regions of the viral genome, into intergenic regions or into a deletion site.

After the introduction of the viral genome and the DNA comprising a host range gene into cells, virus particles comprising the host range gene in the viral genome are isolated/enriched from these cells in a second step. Methods for the isolation/enrichment of viral particles are known to the person skilled in the art. These techniques include e.g. the use of marker genes in the nucleic acid sequence that is introduced into the viral genome. If the marker gene is a selection marker (e.g. a resistance gene) only those recombinant viruses that contain the marker will replicate in infected cells under selective pressure (e.g. if an antibiotic is present). Alternatively or additionally colour markers (e.g. the green fluorescent protein) could be used. If no selection marker is to be used it is possible to isolate and purify recombinant viruses by

limited dilution and/or plaque purification followed by screening of the isolated viruses for the presence of heterologous nucleic acids. Of course these methods may also be combined.

- 5 Methods for obtaining an avipoxvirus comprising a Vaccinia virus host range gene and at least one additional heterologous nucleic acid are known to the person skilled in the art and correspond basically to the method for obtaining the Avipoxvirus according to the present invention as described above. Basically there are three preferred alternatives:
- 10 According to a first alternative a DNA comprising the at least one additional heterologous sequence and an avipoxvirus genome already comprising an vaccinia virus host range gene in the viral genome are introduced into cells in which the virus is able to reproductively replicate. As pointed out above the DNA is preferably a DNA that is
- 15 capable to specifically recombine with the genomic DNA of the avipoxvirus. Then viral particles are isolated/enriched that comprise the at least one additional heterologous sequence in the viral genome from these cells. According to a second alternative a DNA comprising a host range gene as defined above and an avipoxvirus genome already
- 20 harbouring the at least one additional heterologous nucleotide sequence are introduced into cells in which the virus is able to reproductively replicate, wherein the DNA is capable to specifically recombine with the genomic DNA of the avipoxvirus. This is again followed by isolating/enriching virus particles comprising the host range gene in the
- 25 viral genome from these cells. According to the third alternative an avipoxvirus genome and DNA comprising the vaccinia virus host range gene and the additional heterologous nucleic acid sequence are introduced in the cells. The vaccinia virus host range gene and the additional heterologous nucleic acid sequence may be included in one

nucleic acid vector or the host range gene and the heterologous nucleic acid molecule may be included in different nucleic acid vectors. The further steps in the generation of recombinant viruses are as described above.

5

As pointed out above the inventors have shown that the expression of vaccinia virus host range genes in cells productively infected with an avipoxvirus leads to an increase of the avipoxvirus titer produced from the infected cells. According to the above embodiments the expression
10 of the vaccinia virus host range genes was achieved by including functional vaccinia virus host range genes into the viral genome of the avipoxvirus, wherein the host range gene is under the regulation of the natural promoter sequence, any other suitable Vaccinia virus promoter, or any other promoter functional in avipoxvirus infected cells.

15

However, the same results can also be achieved if the functional host range gene is provided by the cell that allows productive replication of the Avipoxvirus. The Vaccinia virus host range gene may be any host range gene as defined above. Preferred host range genes are the
20 Vaccinia virus host range genes for human cells, including the vaccinia virus genes C7L, K1L and E3L. Most preferred is C7L. If not stated otherwise all definitions given above, including the definitions of the viruses, promoters, genes, terms also apply for the following embodiments. Also the order of preferred to most preferred
25 embodiments applies to the following section if not indicated otherwise.

Thus, according to a first alternative of this embodiment the invention concerns a cell comprising a Vaccinia virus host range gene or a homologue of said host range gene, wherein the host range gene is not

part of a Vaccinia virus genome. The invention further concerns these cells infected with an Avipoxvirus, i.e. the invention further concerns cells comprising an Avipoxvirus genome. The Avipoxvirus that is to be used for the infection of the cells or the genome of the Avipoxvirus that is comprised in the cell may or may not comprise a Vaccinia virus host range gene or a homologue thereof in the viral genome. Preferably the cell comprising the Avipoxvirus genome comprises a Vaccinia virus host range gene or homologue thereof, wherein the host range gene or homologue thereof is neither part of a Vaccinia virus genome nor part of the Avipoxvirus genome.

The host range gene is preferably a host range gene or a homologue thereof as defined above, i. e. preferably a host range gene for human cells moer preferably a a host range gene selected from E3L, C7L and K1L.

The host range gene may be integrated in the cellular genome. Methods to generate cell lines containing a foreign gene in the cellular genome are known to the person skilled in the art. According to this embodiment the most preferred cell lines in which the vaccinia virus host range gene is to be stably integrated are avian cell lines (see above), in particular QT35 cells. According to the present invention the Vaccinia virus host range gene comprised in the cellular genome is a functional gene as defined above.

Alternatively, the host range gene may be part of a non-integrated DNA. The non-integrated DNA may be a plasmid DNA that has been introduced into the cell by conventional techniques, before or after the cell is infected with the Avipoxvirus. Moreover, the non-integrated DNA

may be any DNA that persists in the cell without integrating into the cellular genome. Examples for such a persisting, non integrating DNA are recombinant viral genomes, such as Herpesviral genomes and vectors derived from Herpesviral genomes. According to this
5 embodiment the cell may be any cell allowing the productive replication of Avipoxviruses, including primary cells such as CEF cells.

The Avipoxvirus may be any Avipoxvirus as defined above, including recombinant Avipoxviruses.

10

In a second alternative of this embodiment the invention concerns a cell comprising a Vaccinia virus host range gene or a homologue of said host range gene and an Avipoxvirus genome, wherein the host range gene or the homologue of said host range gene is not part of the Avipoxvirus
15 genome. If not indicated otherwise the definitions, the preferred embodiments as well as the order of preferred to most preferred embodiments corresponds to that of the first alternative of this embodiment as shown above. In particular the host range gene may be inserted into the cellular genome or may be part of a non-integrated
20 DNA. However, in addition to the first alternative of this embodiment the second alternative also includes the possibility that the vaccinia virus host range gene or homologue thereof is part of a vaccinia virus genome. Thus, the invention also relates to cells that comprise an Avipoxvirus genome and a Vaccinia virus genome. The Vaccinia virus
25 host range genes are expressed and exert a positive effect on the replication of the Avipoxvirus, resulting in an increased amount of Avipoxvirus produced from said cells compared to cells not comprising a Vaccinia virus genome.

Cells comprising a Vaccinia virus genome as well as an Avipoxvirus genome can be easily obtained by infecting a suitable cell with both, a Vaccinia virus and an Avipoxvirus. If the infected cell allows a productive replication of both, Vaccinia virus and Avipoxvirus, the result of the
5 coinfection is a mixture of both viruses. For most applications it is desirable to obtain an Avipoxvirus preparation without Vaccinia virus contamination. To arrive in such a Vaccinia virus free preparation it is possible either to use specific Vaccinia virus strains that do infect the cells but that do not productively replicate in said cells or to use specific
10 cells or cell lines that allow the reproductive replication of the avipoxvirus but not of the Vaccinia virus.

The above defined cells according to both alternatives of the present invention can be used in a method for amplifying an avipoxvirus
15 characterized in that the cells comprising a Vaccinia virus host range gene or a homologue of said host range gene are infected with the avipoxvirus. The cells are cultivated and the viral particles produced by said cells are isolated/enriched. Alternatively it is possible either to introduce the Avipoxvirus in the cell before introducing the Vaccinia
20 virus host range gene or to introduce the Avipoxvirus and the Vaccinia virus host range gene at the same time. The avipoxvirus may be any poxvirus as defined above, more particularly a wild-type Avipoxvirus, an attenuated Avipoxvirus or a recombinant Avipoxvirus lacking a Vaccinia virus host range gene in the viral genome or a wild-type Avipoxvirus, an
25 attenuated Avipoxvirus or a recombinant Avipoxvirus having a Vaccinia virus host range gene in the viral genome

Moreover the invention concerns the use of a Vaccinia virus host range gene or an homologue thereof to increase the titer of avipoxviruses

produced from avian cells after infection of said cells with said avipoxvirus, wherein the host range gene is expressed in said cells.

- 5 Furthermore the invention relates to a method for increasing the titer of avipoxviruses produced from avian cells by infecting cells comprising a Vaccinia virus host range gene or a homologue of said host range gene with said avipoxvirus.

Summary of the Invention

The invention in its various embodiments may be characterized by the following items

5

1. Avipoxvirus comprising in the viral genome a Vaccinia virus host range gene or a homologue of said host range gene, with the proviso that the host range gene is not the E3L gene if the avipoxvirus is a recombinant canarypoxvirus comprising in the viral genome the Vaccinia virus K3L gene as well as expression cassettes for HIV gag-pro, gp120/TM and a Nef/Pol poly-epitope string, respectively.

10

2. Avipoxvirus according to item 1, wherein the Vaccinia virus host range gene is a host range gene for human cells.

15

3. Avipoxvirus according to anyone of items 1 to 2, wherein the host range gene is selected from the Vaccinia virus genes E3L, C7L and K1L.

20

4. Avipoxvirus according to anyone of items 1 to 3, selected from the group consisting of Fowlpoxvirus and Canarypoxvirus.

25

5. Avipoxvirus according to anyone of the items 1 to 4 comprising in the viral genome at least one additional heterologous nucleic acid sequence.

6. Avipoxvirus according to item 5, wherein the additional heterologous nucleic acid sequence is selected from a sequence coding for at least one antigen, antigenic epitope, and/or a therapeutic compound.

7. Pharmaceutical composition comprising the avipox virus according to anyone of the items 1 to 6 and a pharmaceutically acceptable carrier, diluent and/or additive.

5

8. Vaccine comprising the avipoxvirus according to anyone of the items 1 to 6.

10

9. The virus according to anyone of items 1 to 6, the composition according to item 7 or the vaccine according to item 8 as drug for affecting, preferably inducing, an immunological response in a living animal, including a human.

15

10. Use of the virus according to anyone of the items 1 to 6 for the preparation of a medicament or a vaccine.

20

11. Method for introducing a homologous and/or a heterologous nucleic acid sequence into target cells comprising the infection of the target cells with the virus according to item 5 or 6.

25

12. Method for producing a peptide, protein and/or virus comprising

- a) infection of a host cell with the virus according to anyone of items 1 to 6,
- b) cultivation of the infected host cell under suitable conditions, and
- c) isolation and/or enrichment of the peptide and/or protein expressed from the viral genome and/or of the virus produced by said host cell.

13. Method for affecting, preferably inducing an immunological response in a living animal body including a human comprising administering the virus according to anyone of the items 1 to 6, the composition according to item 7 or the vaccine according to item 8 to the animal or human to be treated.
14. The method according to item 13, wherein the animal is immunocompromised.
15. A cell containing the virus according to any of items 1 to 6.
16. Method for the amplification of an avipoxvirus according to anyone of items 1 to 6, characterized in that
- cells, in particular avian cells, are infected with the avipoxvirus according to anyone of items 1 to 6, wherein the cells are cells allowing the reproductive replication of the avipoxvirus
 - the cells are cultivated and
 - the viral particles produced by said cells are isolated/enriched.
17. Method for obtaining the avipox virus according to anyone of items 1 to 6 comprising the following steps:
- introducing an avipox virus genome that optionally comprises in the viral genome heterologous nucleic acids as defined in anyone of items 5 to 6 and a DNA comprising a host range gene as defined in anyone of items 1 to 3 into cells in which the virus is able to reproductively replicate, wherein the DNA is capable to specifically recombine with the genomic DNA of the avipoxvirus
 - isolating/enriching virus particles comprising the host range gene in the viral genome from these cells.

18. Method for obtaining the avipoxvirus according to anyone of items 5 to 6, comprising the following steps:

- introducing the genome of an avipoxvirus according to anyone of items 1 to 4 and a DNA comprising the at least one additional heterologous sequence into cells in which the virus is able to reproductively replicate, wherein the DNA is capable to specifically recombine with the genomic DNA of the avipoxvirus
- isolating/enriching virus particles comprising the at least one additional heterologous sequence in the viral genome from these cells.

19. Method for obtaining the avipoxvirus according to anyone of items 5 to 6, comprising the following steps:

- introducing an avipox virus genome harbouring the at least one additional nucleotide sequence and a DNA comprising a host range gene as defined in anyone of items 1 to 3 into cells in which the virus is able to reproductively replicate, wherein the DNA is capable to specifically recombine with the genomic DNA of the avipoxvirus
- isolating/enriching virus particles comprising the host range gene in the viral genome from these cells.

20. Cell comprising a Vaccinia virus host range gene or a homologue of said host range gene, wherein the host range gene or the homologue of said host range gene is not part of a Vaccinia virus genome.

21. Cell according to item 20, infected with an avipoxvirus.

22. Cell, in particular a cell according to item 20, comprising a Vaccinia virus host range gene or a homologue of said host range gene and an Avipoxvirus genome, wherein the host range gene or the homologue of said host range gene is not part of the Avipoxvirus genome.
23. Cell according to anyone of items 20 to 22, wherein the cells allow the reproductive replication of the avipoxvirus
24. Cell according to anyone of items 20 to 23, wherein the host range gene is a host range gene as defined in anyone of items 2 to 3.
25. Cell according to anyone of items 20 to 24, wherein the host range gene is integrated in the cellular genome.
26. Cell according to anyone of items 20 to 24, wherein the host range gene is part of a non-integrated DNA.
27. Cell according to anyone of items 20 to 26, namely an avian cell.
28. Cell according to anyone of items 21 to 27, wherein the avipoxvirus genome is a recombinant avipoxvirus genome.
29. Method for amplifying an avipoxvirus characterized in that
- the cells according to anyone of items 21 to 28 are cultivated and
 - that the viral particles produced by said cells are isolated/enriched.

30. Use of a Vaccinia virus host range gene or an homologue thereof, in particular a host range gene as defined in anyone of items 2 to 3 to increase the titer of avipoxviruses produced from avian cells after infection of said cells with said avipoxvirus, wherein the host range gene is expressed in said cells.

5

31. Method for increasing the titer of avipoxviruses produced from avian cells by infecting cells as defined in item 20 with said avipoxvirus or by cultivating cells as defined in anyone of items 21 to 28.

10

Short Description of the Figures

Figure 1: Plasmid map of integration vector pBNCaPVX06

The plasmid contain two regions that are homologous to the Canarypox genome (Flank1, corresponding to the sequence of SEQ. ID: No. 2 and Flank 2, corresponding to the sequence of SEQ. ID: NO. 3). These sequences direct the homologous recombination of the sequences located between Flank1 and Flank 2 into the corresponding location of the viral genome. The integration site into the Canarypox virus genome is located between the TK-gene (Thymidine kinase gene) and a gene named Ca.X with unknown function (see Figure 3). NPTII = neomycin resistance gene (expressed from the PS promoter which is a Vaccinia virus strong synthetic promoter); IRES = internal ribosomal entry site; EGFP = enhanced green fluorescence protein. Bsal= restriction enzyme recognition site for Bsal; prT3= T3 promoter sequence derived from plasmid Bluescript pBSK+ (Stratagene, Inc.); prT7= T7 promoter sequence derived from plasmid Bluescript pBSK+ (Stratagene, Inc.). In this description the designations pBNCaPVX06 and pBNXCaPV06 are used interchangeably and refer to the same plasmid.

Figure 2: Plasmid map of integration vector pBNCaPV08

This plasmid basically corresponds to pBNCaPVX06 explained in the legend to figure 1. pBNCaPV08 additionally comprises the C7L gene derived from MVA (Modified Vaccinia Ankara) expressed from the natural C7L promoter (see Figure 4). The C7L gene from MVA shows the same nucleotide sequence than the C7L gene in Vaccinia virus.

Figure 3: Graphical overview of the intergenic region of the Canarypox genome used for the insertion of Vaccinia virus host

range genes. Ca.6, Ca.5, Ca.X, Ca.3: Canarypox virus genes 6, 5, X and 3, respectively; Ca.TK: Canarypox virus Thymidine kinase gene; Flank1 (SEQ. ID: NO. 2) is a DNA fragment comprising parts of the Ca.6 gene, the entire Ca.5 gene and the entire Ca.Tk gene. Flank2 (SEQ. ID: 3) is a DNA fragment comprising the entire Ca.X gene and parts of the Ca.3 gene.

Figure 4: Sequence of C7L region of MVA (Modified Vaccinia Ankara). This sequence corresponds to SEQ.ID:NO 1.

Figure 5: PCR products for the recombinant Canarypoxviruses canBNX01 and canBN01. Figure 5A: PCR product for canBNX01 shown on a 0.8% agarose gel. lane 1: 100bp marker; lane 2: 1 kb marker; lanes 3-5: different canBNX01 isolates; lane 6: CaPV wildtype; lane 7: pBNCaPVX06; lane 8: water control. **Figure 5B:** PCR product for canBN01 shown on a 0.8% agarose gel. lane 1: 100bp marker; lanes 2-4: different canBN01 isolates; lane 5: water control; lane 6: pBNCaPV08; lane 7: CaPV wildtype

Figure 6: Multistep Growth Curve of recombinant CaPV on Various Cell lines. Amplification of canBNX01 (recombinant CaPV comprising the marker gene cassette but not the Vaccinia virus C7L gene) and canBN01 (recombinant CaPV expressing the C7L gene and the marker gene) in the cell lines BHK-21, Vero, 143B, HaCaT, Hela and MRC-5 and in CEF cells. Virus amplification (fold increase in virus yield above the input level in 6-well-plates) was determined by dividing the virus yield at 96 hours by the input of 5×10^4 (moi 0.05). A ratio of 1,0 means that output = input. The ratios represent the average values of three experiments. Standard errors are indicated by bars.

Figure 7: Nucleotide Sequence of Flank 1 (part A, above) and Flank 2 (part B, below).

Examples

**EXAMPLE 1: Construction of recombinant Canarypox canBNX01
(pS NPTII IRES EGFP) and canBN01 (pS NPTII IRES EGFP C7L-MVA)**

SUMMARY:

This example describes the generation of recombinant Canarypox virus using NPTII (neomycin resistance gene) and EGFP (green fluorescent protein) selection. The Vaccinia Virus host range gene C7L was cloned into an intergenic region of Canarypox by homologous recombination. After two plaque purifications (PP) there was no wild type virus detectable but only recombinant virus. Sequencing of the integration region showed proper integration and no mutations. RT-PCR showed successful expression of the integrated genes, namely the C7L gene from Modified Vaccinia Ankara and the marker gene cassette. The recombinant virus was shown to be stable up to passage number twenty, even without the selective pressure of Geneticin®.

Introduction:

The aim of this example was to construct a recombinant Canarypox virus expressing the Vaccinia Virus host range gene C7L plus marker cassette (canBN01) and a recombinant Canarypox expressing the marker cassette alone (canBNX01). Therefore the integration vectors pBNCaPVX06 (see Figure 1) and pBNXCaPV08 (see Figure 2) had been cloned. Both contain two flanks homologous to the Canarypox virus genome, a marker cassette (NPTII = neomycin resistance, IRES =

internal ribosomal entry site, EGFP = enhanced green fluorescence protein) and for pBNCaPV08 additionally C7L derived from MVA (Modified Vaccinia Ankara) expressed by the natural C7L promoter (see Figure 4; SEQ ID: No 1). C7L from MVA shows the same nucleotide sequence as C7L in Vaccinia virus. The marker cassette is expressed by PS promoter (Vaccinia strong synthetic promoter). The integration site into the Canarypox virus is located in-between the TK-gene (Thymidine kinase) and a gene named Ca.X with unknown function (see Figure 3).

Material:

Recombination vector	pBNCaPVX06 (pBS PS NPTII IRES EGFP) (Fig. 1) pBNCaPV08 (pBS PS NPTII IRES EGFP C7L-MVA) (Fig. 2)
Cells	CEF (Chicken Embryo Fibroblast)
Virus	CaPV crude stock 3.2E+06 TCID ₅₀ /ml
Transfection kit	Effectene (Roche)
DNA-Extraction	Qiagen Blood DNA Kit (Qiagen)
PCR for Wildtype	Taq Polymerase (Roche) Primer:#487: 5'-agcggctttaaattggagatttc-3' Primer: #488: 5'-gttattgttcggaatagaagac-3'
Sequencing	Expand Polymerase (Roche) Big Dye Terminator Kit (PE)
Titration	Anti-CaPV Serum (Guinea Pig #433/1) Anti-Guinea-Pig IgG-POD (Sigma) TMB (Seramun)

Reverse Transcriptase	Rneasy Mini Kit (Qiagen) DNase (Roche) M-MLV RT, RNAsin (Promega) Taq Polymerase (Roche) Primer: #504: 5'-ttactgtacagctcgtccatgc-3' #505: 5'-atgggatcggccattgaacaag-3' #506: 5'-ggcggcggtcacgaactc-3' #498: 5'-ttaatccatggactcataatc-3' #496: 5'-tatacagcacgaattcgacatcatta-3' #497: 5'-ctatacgggattaacggatgttc-3'
-----------------------	---

Method and Results:

5 1. Homologous Recombination (HR):

Plasmids pBNCaPV09 and pBNCaPVX06 (see Figures 1 and 2) were linearised with Restriction Enzyme Bsal at 50°C over night (cuts in the backbone of the plasmids). The digests were purified with a PCR purification kit (Qiagen) and eluted with 50ul H₂O.

10 CEF cells were seeded into 6-well-plates using RPMI 10% FCS. Next day 60 to 80% confluent cells were infected with CaPV crude stock at a moi of 0.1 and 0.01, respectively. Infection was left for one hour at 37°C, virus suspension was removed, and cells were washed with RPMI and 1.6 ml RPMI 2% FCS were added.

Transfection was carried out using Effectene Kit (Roche) following the instruction manual: linearised plasmid (50µl) was mixed with 80µl Buffer EC and 8µl Enhacer were added. The solution was mixed and left for 5 min at room temperature. Then 25µl Effectene was added, mixed and left for 10 min at room temperature. Finally 600µl RPMI 2% FCS were added and transferred onto the cells. The transfection was left for four days at 37°C until fluorescence was detectable. The infection with moi 0.1 seemed to be more effective as clearer fluorescence was detectable. Plates were freeze/thawed three times and finally frozen down at -20°C.

2. Passage and Plaque Purification:

As pointed out above the plasmids used to generate recombinant canarypoxvirus contain a Neomycin resistance gene. Thus, it is possible to select for recombinant viruses by adding Geneticin® at a concentration of 200µg/ml to the cell culture media.

Preparation of crude virus stock canBNX01:

After transfection/infection (see above) the obtained virus containing suspension was given to fresh CEF cells under selective pressure in 6 well plates. The supernatant of cells in which fluorescent plaques were obtained was used for purification of recombinant virus by limited dilution under selection pressure of Geneticin® and by screening for cells/plaques in which the green fluorescent protein was expressed. The generation of recombinant virus was confirmed by PCR screening which also allowed to detect residual Wildtype contamination. The plaque purification steps were repeated until no Wiltype contamination was detectable by PCR. Nearly confluent T25 flasks of CEF cells were infected

with 100µl supernatant of 3 wells in which positive clones for canBNCaPVX01 were detected that were free of Wildtype contamination. The media contained Geneticin®. After three days incubation at 37°C positive fluorescence and CPE was detectable. Flasks were freeze/thawed three times and a crude stock was harvested (P7 = passage number seven). An aliquot of 200µl was taken for PCR analysis. The remaining material was frozen at -20°C. The following three passages were conducted in T25 flasks of CEF cells with and without selection pressure of Geneticin®. The working crude stocks (P11) were prepared in T175 flasks of CEF cells with and without Geneticin®. After these passages the recombinant virus was shown to be stable without Wildtype contamination even in the passages without Geneticin®.

Preparation of crude virus stock canBN01:

Nearly confluent T25 flasks of CEF cells were infected with 100µl supernatant of 3 wells in which positive clones for canBNCaPV01 were detected. The further purification was done in the same manner as described above.

3. PCR Analysis of recombinant Virus and Wildtype

DNA	5.0µl
H2O	2.95µl
x10 buffer	1.0µl
dNTPs	0.2µl
Primer #487	0.4µl
Primer #488	0.4µl
Taq Polymerase	0.05µl

PCR conditions:

94°C, 5min; 94°C, 30sec; 53°C, 30sec; 68°, 3min; 35 cycles; 68°C, 7min; 4°C hold

5

Controls: pBNCaPVX06 and pBNCaPV08 (plasmids used for integration)

DNA from Canarypox (CaPV) (control for Wildtype)

10

Water control

Size of expected PCR-Products:

pBNCaPVX06 and recombinant virus (canBNX01):

2734bp

pBNCaPV08 and recombinant virus (canBNX01): 3461bp

15

CaPV: 436bp

The PCR analysis (Figure 5A and 5B) has clearly demonstrated that recombinant virus was produced and that the purified viruses were free of input wildtype virus.

20

4. Sequencing of inserted genes

Sequencing was conducted with the ABI Prism sequencing machine according to the manufacturers instructions. A PCR product created with primers #487, #488 and Expand Polymerase was used for sequencing. This PCR product includes partly the integration flanks (F1, F1) and entire NPTII IRES EGFP and C7L regions. The expected sequence could be confirmed.

25

5. Titration of crude stocks canBNX01 and canBN01P11

Virus titers were determined in double titrations and the average titer was calculated as follows:

5 canBNX01 plus Geneticin®: 4.9E+06 TCID₅₀/ml
 canBNX01 without Geneticin®: 3.7E+06 TCID₅₀/ml
 canBN01 plus Geneticin®: 1.3E+07 TCID₅₀/ml
 canBN01 without Geneticin®: 2.2E+06 TCID₅₀/ml

10

6. RT-PCR as expression test

6.1 RNA-Preparation

15 CEF cells were seeded into 6-Well-Plates (5 x 10⁵ cells per well DMEM
 10% FCS) and infected the following day with 100µl of canBNX01 and
 canBN01, respectively. Infection was left for two days until fluorescence
 was detectable. RNA-Extraction was conducted with Rneasy Mini Kit
 (Qiagen) according to manufacturers instructions. RNA concentration
 was measured by OD.

20

6.2 DNase digest for RT-PCR

 RNA 25µl
 DNase (RNase-free) 3µl
 10x buffer A (Roche) 5µl
 H₂O (RNase-free) ad 50µl

25

90min at 37°C

Digest was cleaned up using Rneasy Mini Protocol for RNA Clean up and RNA concentration was measured by OD.

6.3 Reverse Transcriptase

- 5 RNA and Primer #504 (for canBNX08) or #498 (for canBN01) were mixed in a ratio of 2µg RNA to 1µg primer. Water (RNase-free) was added up to a total volume of 10µl. The mixture was left for 5 min at 70°C and then it was incubated on ice.

The following was added:

10	5x buffer	5µl
	dNTP	5µl
	Rnasin	0.5µl
	M-MLV RT	2µl
	H ₂ O (RNase-free)	2.5µl
15	60min 42°C	

RT was cleaned up using PCR purification Kit (Qiagen).

6.4 PCR (Taq Roche)

20	DNA	5µl
	x10 buffer	5µl
	dNTP	1µl
	Primer 1	2µl
	Primer 2	2µl
25	Taq (Roche)	1µl

PCR conditions:

94°C, 5min; 94°C, 30sec; 58°C, 30sec; 68°C, 2min 30sec; 30 cycles;
68°C, 7min; 4°C hold

Samples: RNA before RT-PCR (to detect contaminant DNA)

5 RNA after RT-PCR and clean up

Control: pBNCaPVX06 and pBNCaPV08 (positive controls)

Primer: #505, #506 for canBNX01
#496, #497 for canBN01

Size of expected PCR-Products:

10 pBNCaPVX06 and recombinant virus (canBNX01):
2188bp
pBNCaPV08 and recombinant virus (canBN01):
428bp

15 The expression of the inserted genes could be confirmed as positive by
RT-PCR.

Conclusion:

20 With the described method it was possible to construct a recombinant
Canarypox virus comprising the Vaccinia virus C7L gene. The C7L gene
is derived from MVA (Modified Vaccinia Ankara) and shows the same
nucleotide sequence as the C7L gene in Vaccinia Virus copenhagen.
The selection method was shown to be very effective since there was no
Wildtype virus detectable after two plaque purifications.

25 Sequencing of the inserted genes and parts of the surrounding flanks
showed no mutations that affect the function. RT-PCR showed the
expression of the Vaccinia host range gene C7L under regulation of the
natural promoter.

EXAMPLE 2: Multistep growth curve analysis of recombinant Canarypox canBNX01 (pS NPTII IRES EGFP) and canBN01 (pS NPTII IRES EGFP C7L-MVA)

5

SUMMARY:

The aim of this example was to investigate replication of a recombinant Canarypox virus expressing the human (tissue culture) host range gene C7L under regulation of the natural promoter in a multistep growth curve on several cell lines. Multistep growth curve means that infection is performed on a low moi (multiplicity of infection) which enables to investigate viral spread and replication. The results indicate that the recombinant Canarypox has improved growth properties on CEF cells resulting in titers which are one log higher than those of the control virus expressing the marker cassette only. The replication properties on several mammalian cell lines (human, monkey and rabbit cell lines) remains non-effected, which means that the recombinant virus seems to be as attenuated as the control virus.

20

Introduction:

This example evaluates the growth potential of recombinant Canarypox virus expressing C7L (canBN01; for cloning details see example 1) and the control virus not expressing the C7L gene (canBN01; see example 1) in different cell lines or primary cells. The cell lines/cells used are cell that are permissive for the canarypoxvirus, such as CEF cells and cell lines that are non-permissive for the canarypoxvirus, such as BHK-21, Vero, 143B, HaCaT, Hela, MRC-5 and RK-13 cells. Canarypox is known to be strictly restricted to grow only in avian cells as represented by the

25

primary CEF cells (Esposito et al, 1991; Plotkin et al, 1995, Taylor et al, 1995). Unfortunately, the viral titers are relatively low when compared to other poxviruses, for example MVA. Therefore, the growth properties of a recombinant Canarypox virus expressing the human Vaccinia Virus host range gene C7L (Perkus et al, 1990; Oguiura et al, 1993) were evaluated in CEF cells and it was checked whether the recombinant virus is still not capable replicate on mammalian cell lines.

10 **Material:**

Cell lines: CEF: Chicken Embryo Fibroblast, primary cells
 BHK-21: Baby Hamster Kidney cells, fibroblast cell line
 Vero: African green monkey kidney cells, fibroblast cell line
 15 143B: human Osteosarcoma cell line, TK
 HaCaT: human keratinocyte cell line
 Hela: human cervix carcinoma cell line, epithelial
 MRC-5: human lung cell line, fibroblast
 RK-13: rabbit kidney cell line, epithelial
 20 All cells are cultured in DMEM 10%FCS

Virus: can BNX01, recombinant Canarypox virus expressing marker cassette only (NPTII, IRES, EGFP; regulated by pS strong synthetic promoter)
 25 canBN01, recombinant Canarypox virus expressing marker cassette (NPTII, IRES, EGFP; regulated by pS strong synthetic promoter) and C7L from MVA regulated by the natural promoter

Cell Culture Medium: DMEM plus 2% FCS
DMEM, Gibco
FCS, PAA

5 Other Reagents: RPMI, Gibco
Antibiotic-Antimycotic, Gibco,
PBS, Gibco
Trypsin EDTA (1x), Gibco
Fixing solution: Aceton/Methanol 1:1
10 Incubation solution: PBS plus 3% FCS
Anti-CaPV Serum (Guinea Pig #433/1)
Anti-Guinea-Pig IgG-POD (Sigma)
Staining solution: PBS plus TMB (Seramun) 1:1

15

Methods:

Infection of various cell lines in six-well-plates

20 Each cell line was grown to nearly confluency in three six-well tissue
culture dishes for each of both viruses. The cell monolayers were
infected at a moi (multiplicity of infection) of approximately 0.05 using a
total of 5×10^4 TCID₅₀/ml in 500µl of DMEM for each well. Infection was
left for one hour at 37°C, then cells were washed two times with DMEM
25 to remove unadsorbed virus and incubated with 1000µl DMEM 2% FCS
for four days at 37°C 5% CO₂.

After the infection the cells were scraped into the medium and cells plus medium were freeze-thawed three times to release the viruses from the cells. These viral extracts were titered on CEF cells.

5 Titration of CaPV (Immunostaining with a Canarypox virus specific antiserum):

Titration was performed on CEF cells. Briefly, test cells (CEF) were seeded on 96-well-plates in RPMI 1% Antibiotic/Antimycotic 7% FCS at
10 a concentration of 1×10^4 cells/well and incubated over night at 37°C 5% CO₂. The test samples had already been frozen/thawed 3 times; dilutions of 10^{-1} to 10^{-12} were prepared using RPMI medium. Virus dilutions were distributed onto test cells and incubated for five days at 37°C 5% CO₂ to allow CPE development. Test cells were fixed for 10
15 min, washed with PBS and incubated with polyclonal Canarypox specific antiserum at a 1:1000 dilution in incubation buffer for one hour at RT. After washing twice with PBS the HPR-coupled anti-Guinea-Pig antibody was added at a 1:1000 dilution in incubation buffer for one hour at RT. Cells were again washed twice with PBS and incubated with staining
20 solution until blue spots were visible (15 min). Staining solution was removed and cells were washed with PBS. Every well showing a brown spot was marked as positive for CPE and titre was calculated using the formula of Kaerber (TCID₅₀ based assay) (Kaerber, G. 1931. Arch. Exp. Pathol. Pharmacol. 162, 480).

25

Results:

Recombinant Canarypox virus expressing the Vaccinia Virus host range gene C7L was used to infect triplicate sets of CEF, BHL-21, Vero, 143B, HaCaT, Hela, MRC-5 and RK-13 cells at a low multiplicity of infection (moi 0.05). After infection the virus inoculum was removed and cells were washed two times to remove any unabsorbed free virus particle. Then, infections were left for 4 days; virus extracts were prepared and titrated on CEF cells. Figure 6 plots the ratios of Output/Input for the 6-well plates (Output means total virus production after four days, and Input means the amount of virus used for the initial infections). These ratios give a clear indication of the extent of viral amplification in the various cell types.

As clearly visible, the recombinant Canarypox expressing C7L (canBN01) shows on CEF cells approximately 10 fold higher titers than the control virus (canBNX01). This means an enhancement of about one log for the titer.

When compared to the level of amplification that occurred in all of the mammalian cell lines tested (100 to 1000 fold decreases above input), canBN01 does seem to be severely growth restricted in the cell lines tested. The expression on a Vaccinia Virus human host range gene does not seem to affect replication of Canarypox on mammalian cell lines. Figure 6 clearly demonstrates that amplification by cell to cell spread of canBN01 in the cell lines tested can not be detected.

Conclusion:

The genetic engineering of Canarypox resulting in expression of the Vaccinia Virus human host range gene C7L under regulation of its natural promoter (derived from MVA; Modified Vaccinia Ankara) is

useful to increase viral titers on CEF cells. Canarypox is known to grow to relatively low titers and to grow slower than the other poxviruses (for example MVA). Therefore, Vaccinia virus host range genes are a good tool for increasing production of Canarypox without affecting the attenuated replication properties on a range of mammalian cells.

References for Example 2:

Esposito, J. J. et al.: *Fifth report of the International Committee on Taxonomy of Viruses.* Arch. Virol. Suppl. (1991) 2, 79-102.

Oguirua, N. et al.: *Detection of a protein encoded by the vaccinia virus C7L open reading frame and study of its effect on virus multiplication in different cell lines.* Journal of General Virology (1993) 74, 1409-1413.

Perkus, M. et al.: *Vaccinia Virus Host Range Genes.* Virology (1990) 179, 276-286.

Plotkin, S. A. et al.: *The Safety and Use of Canarypox Vectored Vaccines.* Dev Biol Stand. Basel, Karger (1995) Vol. 84, 165-170.

Taylor, J. et al.: *Biological and immunogenic properties of a canarypox.rabies recombinant, ALVAC-RG (vCP65) in non-avian species.* Vaccine (1995) Vol. 13, No. 6, 439-549.

Claims

5

1. Avipoxvirus comprising in the viral genome a Vaccinia virus host range gene or a homologue of said host range gene, with the proviso that the host range gene is not the E3L gene if the avipoxvirus is a recombinant canarypoxvirus comprising in the viral genome the Vaccinia virus K3L gene as well as expression cassettes for HIV gag-pro, gp120/TM and a Nef/Pol poly-epitope string, respectively.
2. Avipoxvirus according to claim 1, wherein the host range gene is selected from the Vaccinia virus genes E3L, C7L and K1L.
3. Avipoxvirus according to anyone of the claims 1 to 2 comprising in the viral genome at least one additional heterologous nucleic acid sequence, preferably selected from a sequence coding for at least one antigen, antigenic epitope, and/or a therapeutic compound.
4. Pharmaceutical composition or vaccine comprising the avipox virus according to anyone of the claims 1 to 3 and a pharmaceutically acceptable carrier, diluent and/or additive.
5. Method for the amplification of an avipoxvirus according to anyone of claims 1 to 3, characterized in that
 - cells, in particular avian cells, are infected with the avipoxvirus according to anyone of claims 1 to 3, wherein the cells are cells allowing the reproductive replication of the avipoxvirus

- the cells are cultivated and
- the viral particles produced by said cells are isolated/enriched.

5 6. Cell comprising a Vaccinia virus host range gene or a homologue of said host range gene, wherein the host range gene or the homologue of said host range gene is not part of a Vaccinia virus genome and wherein the cell allows a reproductive replication of Avipoxvirus.

10 7. Cell according to claim 6, infected with an avipoxvirus.

15 8. Cell, in particular a cell according to claim 6, comprising a Vaccinia virus host range gene or a homologue of said host range gene and an Avipoxvirus genome, wherein the host range gene or the homologue of said host range gene is not part of the Avipoxvirus genome and wherein the cell allows a reproductive replication of the Avipoxvirus.

20 9. Use of a Vaccinia virus host range gene or an homologue thereof, in particular a host range gene as defined in claim 2 to increase the titer of avipoxviruses produced from avian cells after infection of said cells with said avipoxvirus, wherein the host range gene is expressed in said cells.

25 10. Method for increasing the titer of avipoxviruses produced from avian cells by infecting cells as defined in claim 6 with said avipoxvirus or by cultivating cells as defined in anyone of claims 7 to 8.

Abstract


The invention concerns an Avipoxvirus comprising in the viral genome a Vaccinia virus host range gene or a homologue of said host range gene.

5 The invention further relates to cells, preferably avian cells, comprising a Vaccinia virus host range gene or a homologue of said host range gene. Moreover the invention concerns the use of a Vaccinia virus host range gene or an homologue thereof to increase the titer of avipoxviruses produced from cells after infection of said cells with the avipoxvirus,

10 wherein the host range gene is expressed in said cells.


1/7

Fig. 1




2/7

Fig. 2



3/7

Fig. 3



5/7

Modtaget PVS
- 7 AUG. 2002

Figure 5A:





Figure 5B:



- 7 AUG. 2002

Figure 6:




Figure 7A:

ATACTATTCTTCACGGTACATTTAAAAAAGGAATATAGTCAGAAACAGGAAATATACT
TTCACATAACATGGTCTAATTTTGAATGTCCGACGTTAGGAGACGTTAAGTCTTCTTC
ACCTAATACCTGTAATAGAGTAGTTTTAGACGGTAGTAGATACGTTACAAAACCTTTA
ATGATACAATATAAATGGAACCTAACTAGAGAAACGCTGATATTTGTAGGCATTACTGTA
CTAGTAGTAGTAATGATCATATCTGGTTTCTCCTAATATTGCGATTGATACCTGGTGT
ATATTCATCAGTTATTAGATCGTCGTTTCGTAGGAGGGAAAATATTAAGATTTATGGAGG
TATTCTCTACTGTTATGTTTATACCATCATTAGTAATACTTTATACAGCATATATAAGG
AAATCTAAAGTGAAAAATAACTAAATATTATAGTATTTGTAATAAATGGCTACTGGAGA
GATTCGTCTTATTATAGGGCCATGTTTTTCAGGTAAAACAACAGAATTAGTTAGATTAA
TAAGAAGATTTATGATATCGGGACGTAAATGTATAATAATAAACATTGTAGTGATTCC
CGTTATACCGAAGGAGATTTAGAAGCTATATATACTCATGATAAAATTTTCGATGGAAGC
ACTATCGTGTAGCAAATTATTACCTTTAATACCTAAAATTGATAACTTTGAAGTAATAG
GTATAGACGAAGGACAGTTTTTTTGAAGATATAGTAGAATTTAGTGAGATTATGGCTAAT
AAGGGTAAACTGTAAATCATAGCGGCTTTAATGGAGATTTCAAACGACAATTATTTGG
AAACATATTTAACTATTATCTTTATCAGAATCAGTTACTAGTTTAACTGCTATTTGTG
CAGTTTGTA AAAACGAAGCATCTTTTTCTAAGCGCATGACTGATGATAAAGATGTAAAA
GTTATAGGAGGTAAAGAAATGTATACTGCTGTTTGTAGAAAATGCTTTTTTATGAGTC

Figure 7B:

TAATATACGTAATACTTGTACGTACAACCTATGTTAGAATAATTTGCTTAGTATAG
TATATAACAAGTATGTAAAAATAAAATTGATATAAAAGTAGTCTTCTATTCCGAACA
ATAACTATACAAAATGGATTTAGATATTAAATCTTGCAGAAGTATTTACAAAATATGGG
ATAAATATCATTTTTATGACAGGGTATAAATATAAAAATGATAAACAGAGATTTAAAATT
ACAATTTACTGTAAATGTGATTGTTCTATCAAAGAATATCCTTATAGATTTGTTACTGA
GAACTGCTTTTAATGTATATTATTAATAAGTTTAGAGGAAAGTATCTAATCAAATTA
GGATAGAACCCATAGTTAAAAATTAAATCATATATCAATACATGTCAGTTTTTTATCGA
AAAATGGATTTATAAATAAAATGAAAAATAACTTGAATGAAGGAAAAATAACCATGAG
TAAAAAACAGTAAGACGGTCCAGCGTAGACGTGGAACGATGAGGATAATAAGTTTA
CTTGATCCAAGCGCTAGAACATGCAAAAAGCTTATGTACTAAAAATAATAAATAGTT
AAATCTGTTAACTATCACAATCTCTCTTTAAGTCATCTAACAATATTTCTGTGATATT
AGAACCAGAATATAAAGACAAATTAGTGACTCCTCTTATTATTGTAGAAGGTGAAGGAA
AAATATACCATAATAAGAATGATAGTTTTAATCGTGAAGAACCGTATTTTCTAAAAATA
CGACCTACGTTAATGAATCCTATATTATATCAGATTATGGAATGCATTTATAGAGATCT
CAATTATTTGGATCCCGAGAATACGATGGATGAAAAACATTTAAAGATTGTCATCTGT
ATATTAACGGAAATAGGATTATGTCCGCCGACGTAAAATATTTGAAGAATGGTAAACCT
GTAGGAGAAAAATTATCCGTATCCAAGGAAATAGATAAACTGGTTAAAAAAGATCCACA

BN-C7L.ST25.txt
SEQUENCE LISTING

Modtaget PVS
- 7 AUG. 2002

<110> Bavarian Nordic A/S

<120> vaccinia virus host range genes to increase the titer of Avipoxviruses

<130> BN48DK-Avi

<160> 3

<170> PatentIn version 3.1

<210> 1

<211> 615

<212> DNA

<213> artificial sequence

<220>

<223> 1-162 = estimated promoter sequence for C7L in MVA

<220>

<221> promoter sequence C7L

<222> (1)..(162)

<223>

<220>

<221> coding sequence

<222> (163)..(615)

<223>

<400> 1
attaataaac tttaagacat gtgtgttata ctaagatggt tggcttattc catagtagct 60
tgttgaattt ataaacttat gatagtaaaa ctagtaccba atatgtaaag atgaaaaagt 120
aaattactat taacgccgtc ggtattcgtt catccattca gtatgggtat acagcacgaa 180
ttcgacatca ttattaatgg agatatcgcg ttgagaaatt tacagttaca taaaggggat 240
aactacggat gcaaactaaa aattatttcg aatgattaca agaaattaaa gtttagattc 300

BN-C7L.ST25.txt

attatacgcc cagattggtc ggaaatcgac gaggtcaaag gattaaccgt atttgcaaac	360
aactatgcgg tgaaagttaa taaggtagat gacacgttct attacgtaat atatgaggct	420
gtaatacatc tgtataacaa aaaaacagag atattgattt attctgatga tgagaacgaa	480
ctctttaaac actattaccc atacatcagt ctaaatatga ttagtaaaaa gtataaagtt	540
aaagaagaaa actactcatc cccgtatata gaacatccgt taatcccgt tagagattat	600
gagtcctgg attaa	615

<210> 2

<211> 1001

<212> DNA

<213> artificial sequence

<400> 2	
atactattct tcacggtaca tttaaaaaaaa ggaatatagt cagaaacagg aaatatactt	60
tcactataac atgggtcta ttcgaatgtc cgacgttagg agacgttaag tcttcttcac	120
ctaatacctg taatagagta gtttagacg gtagtagata cgttacaaaa acctttaatg	180
atacaatata aatgggaacta actagagaaa cgctgatatt tgtaggcatt actgtactag	240
tagtagtaat gatcatatct ggtttctcac taatattgag attgatacct ggtgtatatt	300
catcagttat tagatcgctg ttcgtaggag ggaaaatatt aagatttatg gaggtattct	360
ctactgttat gtttatacca tcattagtaa tactttatac agcatatata aggaaatcta	420
aagtgaaaaa taactaaata ttatagtatt tgtaataaat ggctactgga gagattcgct	480
ttattatagg gcctatgttt tcaggtaaaa caacagaatt agttagatta ataagaagat	540
ttatgatatc gggacgtaaa tgtataataa taaaacattg tagtgattcc cgttataccg	600
aaggagattt agaagctata tatactcatg ataaaatttc gatggaagca ctatcgtgta	660
gcaaattatt acctttaata cctaaaattg ataactttga agtaatagggt atagacgaag	720
gacagttttt tgaagatata gtagaattta gtgagattat ggctaataag ggtaaaactg	780
taatcatagc ggcttttaaat ggagatttca aacgacaatt atttggaac atattttaaac	840
tattatcttt atcagaatca gttactagtt taactgctat ttgtgcagtt tgtaaaaacg	900
aagcatcttt ttctaagcgc atgactgatg ataaagatgt aaaagttata ggaggtaaag	960
aaatgtatac tgctgtttgt agaaaatgct ttttatgagt c	1001

<210> 3

<211> 1003

<212> DNA

<213> artificial sequence

```

<400> 3
taatatacgt actaaatact tgtacgtaca actatgttag aataatttgc ttagtatagt 60
atataaacia gtatgtaaaa aataaaattg atataaaaagt agtcttctat tccgaacaat 120
aactatacaa aatggattta gatattaaat cttgcagaag tatttaciaa atatgggata 180
aatatcattt tatgacaggg tataaatata aaaatgataa acagagattt aaaattacia 240
tttactgtaa atgtgattgt tctatcaaag aatatcctta tagatttggt actgagaaac 300
tgcttttaat gtatattatt aataagttta gaggaagta tctaatacaa attaggatag 360
aaccatagt taaaaattaa atcatatatc aatacatgtc agttttttat cgaaaaatgg 420
atttataaat aaaatgaaaa ataacttgaa tgaaggaaaa aataaccatg agtaaaaaac 480
cagtaaagac ggtccagcgt agacgtggaa acgatgagga taataagttt acttgtatcc 540
aagcgctaga acatgcaaaa agcttatgta ctaaaaataa taaaatagtt aaatctgtta 600
aactatcaca atctctcttt aagtcattta acaatatttc tgtgatatta gaaccagaat 660
ataaagacia attagtgact cctcttatta ttgtagaagg tgaaggaaaa atataccata 720
ataagaatga tagttttaat cgtgaagaac cgtattttct aaaaatacga cctacgttaa 780
tgaatcctat attatatcag attatggaat gcatttatag agatctcaat tatttggatc 840
ccgagaatac gatggatgaa aaaacattta aagattgtca tctgtatatt aacggaaata 900
ggattatgtc cgccgacgta aaatatttga agaatggtaa acctgtagga gaaaaattat 960
ccgtatccaa ggaaatagat aaactgggta aaaaagatcc aca 1003

```

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.